Short communication

Rainfall and runoff pattern of Coorg district of Karnataka state

H.L. GURUPRASANNA, B.T. PANDURANGA RAVINDRABABU and M.B. RAJEGOWDA

University of Agricultural Sciences, GKVK, Bangalore, Karnataka-560 065.

Cauvery is a major river that provides water both for irrigation and other domestic purposes to Karnataka and its east neighboring states. Its origin is in Coorg District of Karnataka State, Coorg District occupies major portion of catchments area to feed the Cauvery River. Quantum of rainfall received at Coorg District decides the irrigation fate in the State and its neighboring states. It also helps in supplying the water required for domestic purposes in addition to the production of Hydroelectricity. The district falls under hilly Agroclimatic zones of the State and has three taluks i.e., Madikeri, Somwarpet and Virajpet. Coorg district of Karnataka being the place of origin of Cauvery, occupies a major portion of catchments area. In this context a study on rainfall in Coorg district i.e. annual rainfall, seasonal rainfall, rainfall trend during 1951-2002, PET, soil water storage, runoff and amount of water available for the evapotranspiration has been worked out

The monthly rainfall data for the period from 1951 to 2002 has been analyzed making into two groups, a). The normal annual rainfall for 30 years period has been worked out and it is 2725 mm. and b). For the remaining period from 1981 to 2002 the

mean has been worked out and it is 2645 mm and the same has been compared with that of previous normal. The annual rainfall data has been divided into pre-monsoon (January to May), South-West monsoon (June to September) and Northeast monsoon (October to December) seasons and their changes between the two groups has been studied. The annual rainfall trend has been worked out for the period from 1951 to 2002.

It is observed that the pre-monsoon and south west monsoons have been increased in the district and slight increase in the north east monsoon. Mean seasonal and annual rainfall in different seasons and their statistics have been worked out for the period from 1981 to 2002 and provided in Table 1. The average rainfall in the district ranges from 2167 mm in Somwarpet taluk to 3264 mm in Madikeri taluk. The rainfall decreases from South-Eastern part of the district towards North and North-Western part.

Table 1 reveals that Coorg district receives an average annual rainfall of 2645 mm (1981-2002) against the previous normal of 2725 mm (1951-1980). During the recent period, Pre-monsoon rainfall contributes about 233 mm (8.8 % of the

Table 1: Seasonal and annual mean rainfall (mm) during different seasons and their statistics for the period from 1981 to 2002.

Taluks	Statistical parameters	Pre-monsoon (Jan May)	South-West monsoon (June – Sept.)	N -E monsoon (OctDec.)	Annual Total
Madikeri Somwarpet	Mean rainfall	310	2649	305	3264
	S.D	118	595	120	629
	C.V. %	45	22	40	19
Somwarpet	Mean rainfall	217	1654	296	2167
	S.D.	117	410	110	449
	C.V. %	54	110,000	42	21
Virajpet	Mean rainfall	172	2012	320	2504
	S.D.	143	546	131	580
	C.V. %	53	27	43	23
District	Mean(1981-2002) Nor.(1951-1980)	233 246	2105 2175	307 304	2645 2725

Table 2: Comparison of recent (1981-2002) deficit rainfall years (per cent) with the Normal deficit rainfall years (1951-1980)

Taluks	Pre-Monsoon		S-W monsoon		N-E monsoon		Annual rainfall	
	Normal	Recent years	Normal	Recent years	Normal	Recent years	Normal	Recent years
Madikeri	37	40	18	20	38	35	14	20
Somwarpet	42	50	24	30	38	50	19	25
Virajpet	43	65	24	40	43	50	20	35
District	41	52	22	30	40	45	18	27

annual) and South-West rainfall contributes about 2105 mm (79.6 % of the annual) and rest of 307 mm (11.6 % of the annual) from the North-East rainfall. Madikeri taluk receives highest annual rainfall of 3264 mm rainfall in the district. Realizing the decrease in rainfall during recent years, the deficit years for the entire period have been worked cut and compared with the normal deficit years (Table 2).

It is observed that in all the taluks, recent 22 years data show more number of deficit years. This increase in deficit years is more particularly during pre-monsoon and northeast monsoon period. The annual rainfall trend indicates higher deficit years in all the taluks.

Water balance

Monthly climatic water balance has

Table 3: Mean monthly water balance for the Coorg district

Month	Rainfall (mm)	Infiltration (mm)	Net rain fall (mm)	PET (mm)	AET (mm)	Run off (mm)
						0.0
Jan.	5.8	0.0	5.8	92.2	46.2	0.0
Feb.	6.3	0.0	6.3	96.9	30.0	0.0
Mar.	21.0	0.0	21.0	121.7	34.9	0.0
Арг.	76.4	0.0	76.4	129.6	80.7	0.0
May.	123.9	6.2	117.7	143.4	119.3	0.0
Jun.	496.0	24.8	471.2	100.0	100.0	229.8
Jul.	896.1	44.8	851.3	86.0	86.0	765.3
Aug.	498.8	24.9	473.9	92.2	92.2	381.7
Sep.	214.5	10.7	203.8	86.8	86.8	117.0
Oct.	201.5	10.1	191.5	82.9	82.9	108.5
Nov.	87.4	0.0	87.4	83.7	83.7	8.3
Dec.	18.4	0.0	18.4	86.0	71.2	0.0
Total	2646	121.5	2524.6	1201.4	913.9	1610.6

been worked out using the Thornthwaite and Mather (1955) book keeping procedure. The Soil depth, water holding capacity, infiltration rate and available water capacity values (150 mm for every 150 cm depth of the soil at its field capacity) have been taken from the publication of Shivaprasad et al (1998).

The mean monthly PET values have been taken from Khambete and Biswas, (1992). The mean monthly infiltration of about 4.5 per cent (Shivaprasad et al 1998), has been worked out and net rainfall has been obtained. The excess rainwater after saturating the 150 cm of soil goes as runoff and is given in the last column.

From the Table 3, it is observed that, Coorg district receives the highest rainfall of 896.1 mm in the month of July followed by August (498.8) and June and 496.0 mm months. About 4.6 per cent (121.5 mm) of the normal annual rainfall infiltrates beyond root zone and adds to the ground water recharge leading to net rainfall of 2524.6 mm. Annual Potential Evapotranspiration is about 1201 mm and 1610.5 mm will go as annual runoff. Runoff is found only during the period from June to October months. Hence the rainfall during this period has greater impact on agriculture in the Cauvery basin as it contributes much to the reservoir. There is adequate soil moisture storage through rainwater to meet the atmospheric demand during June, July, August, September, October and November months.

REFERENCES

Khambete, N.N and Biswas, B.C. 1992. Weekly Potential Evapotranspiration over India, Meteorological Monograph, Agromet/No.14/1992.

Shivaprasad, C.R., Reddy, R.S, Seghal, J and Velayudam, M. 1998. Soils of Karnataka for Optimizing Land Use. NBSS Publication 47 b (Soils of India series), National Bearou Soil Survey and Land Use Planning, Nagapur, P111.