

Rainfall analysis and crop planning in lower Shiwalik foothills of Punjab

PARVENDER SHEORAN, SOM PAL SINGH¹ and VIRENDER SARDANA

Oilseeds Section, Department of Plant Breeding, Genetics and Biotechnology, ¹Department of Agril. Meteorology, PAU, Ludhiana-141 004, Punjab
Email : sheoran76@rediffmail.com

ABSTRACT

Daily rainfall data of 21 years (1984-2004) recorded at RRSKA, Ballawal Saunkhri, District Nawanshahr, Punjab was examined for long term averages of annual, seasonal, monthly and weekly rainfall and its temporal variability. Coefficient of variation of 27.1 per cent indicated that the annual rainfall was more or less stable over the years. The season-wise per cent contribution to annual rainfall was 7.2, 13.2 and 79.6 per cent of summer, *rabi* and *kharif* seasons, respectively. With in the rainy season, August was the highest rainfall contributing month (35.8%) followed by July (35.4%). Mean weekly precipitation amount and its assurance reaches the peak (>50 mm/week) during 27th SMW to 36th SMW and again declined thereafter. The earliest onset of rainy season occurred in 24th SMW. The normal onset of rainy season was observed as 26th SMW with CV of 5.2 per cent. There is an ample scope for rain water harvesting from July to September which can be utilized as crop saving irrigation as well as pre-sowing irrigation for succeeding *rabi* crops which are generally sown on residual soil moisture.

Key words: Rainfall pattern, probability analysis, crop planning

Agriculture being mainly rainfed, the lower shiwalik foothills of Punjab state, India locally known as 'Kandi' is characterized by undulating to escapement topography, uneven and erratic distribution of rainfall, deep ground water table, severe soil erosion, inherent poor soil fertility and frequent crop failures. Since rainfall is the only source of moisture, the spatio-temporal distribution of rains holds the key in determining the fate of entire crop productivity in the region. Knowledge of average monthly, seasonal and annual rainfall is helpful in understanding the general picture of the particular region but the weekly rainfall data analysis gives more useful and precise informations for the rainfall based crop planning (Tiwari *et al.*, 1992). Rainfall probabilities (wet and dry spells) could be made use of in selection of different cropping systems (Jadhav *et al.*, 1999). The coincidence of wet spells with the sensitive phenophases sometimes may be more detrimental to the crop development. On the other hand, the occurrence of dry spell at the time of ripening would become beneficial. Therefore, the probabilities of wet and dry spells can serve as a basic input for establishing precise crop-weather relationships to take some useful decisions for crop management practices, contingent crop planning and related farm operations for sustaining crop production in the area. Such an analysis has also been reported

from other parts of the country to follow a profitable crop planning system under rainfed conditions (Sarma *et. al.*, 1996 and Dingre and Habib, 2006). Hence, an attempt has been made to analyze the daily rainfall data of Ballawal Saunkhri for evolving rainfall based cropping system.

MATERIALS AND METHODS

The daily rainfall data for 21 years (1984-2004) recorded at Agro-meteorological Observatory (31°6'5" N, 76°27'26" E and 355 m above mean sea level), PAU Regional Research Station for *Kandi* Area (RRSKA) of P.A.U Ludhiana, Ballawal Saunkhri, District Nawanshahr, Punjab is examined for analyzing rainfall distribution pattern of the area, probability of wet and dry spells and its implications for crop planning. An attempt has been made to use a standard week in drought analysis using Markov Chain Model and planning crops accordingly. A week receiving less than 21 mm rainfall is taken as dry spell week and a week receiving 21 mm or more rainfall as a wet week (Subramaniam and Raju, 1988). From the daily rainfall data, annual, seasonal, monthly and weekly total rainfall was worked out for each year.

The probability of occurrence of a dry week {p(d)}, two consecutive dry weeks {p(dd)} and three




Fig. 1: Mean weekly rainfall distribution at Ballowal Saunkhri (1984-2004)




Fig. 2: Mean monthly probabilities of dry and wet spell sequences

consecutive dry weeks {p(ddd)} was computed by Markov Chain process (Robertson, 1976). Similarly, p(w), p(ww) and p(www) were calculated in the same way and the notations used are having the same meaning except the week being wet.

RESULTS AND DISCUSSION

Annual rainfall

The compilation of 21 years daily rainfall data of RRSKA, Ballowal Saunkhri showed that the average annual rainfall of the station was 1129.2 mm spread

over 52 rainy days with 27.1 per cent coefficient of variation (CV) indicating low variability over the years (Table 1). The rainfall was above normal in 8 years and below normal in 13 years. The maximum and minimum rainfall of 2041.1 and 816.5 mm was recorded in the years 1988 and 2002 on 69 and 43 rainy days, respectively.

Seasonal rainfall

The seasonal rainfall distribution (Table 1) in terms of summer (March to May), *kharif* (June to September) and *Rabi* (October to February) seasons

Table 1: Mean monthly and seasonal rainfall (mm) at Ballawal Saunkhri (1984-2004)

Months	Rainfall (mm)	Contribution to total (%)	STD	SE±	CV (%)	Rainy days
March	26.0	2.3	17.9	3.9	68.6	2.8
April	20.4	1.8	18.7	4.1	91.7	2.3
May	34.9	3.1	33.7	7.4	96.7	2.9
<i>Summer season</i>	81.3	7.2	41.9	9.1	51.5	8.0
June	94.5	8.4	52.4	11.4	55.4	5.4
July	318.9	28.2	174.1	38	54.6	11.7
August	321.6	28.5	117.1	25.6	36.4	11.9
September	164.2	14.5	143.5	31.3	87.4	5.8
<i>Kharif season</i>	899.2	79.6	296.6	64.7	33.0	34.7
October	33.1	2.9	57.5	12.5	173.7	1.4
November	8.0	0.7	14.7	3.2	184.5	1.0
December	30.0	2.7	48.5	10.6	161.9	1.5
January	36.8	3.3	33	7.2	89.8	2.5
February	40.9	3.6	32.7	7.1	80.0	3.2
<i>Rabi season</i>	148.8	13.2	87.8	19.2	59.1	9.7
Grand total	1129.2	100.0	305.7	66.7	27.1	52.3

revealed that the region enjoys a well defined pattern of seasonal rainfall. The rainfall received was the highest in *kharif* season (899.2 mm) followed by *rabi* with 148.8 mm. The season-wise per cent contribution to total annual average rainfall was 7.2, 13.2 and 79.6 per cent of summer, *rabi* and *kharif* seasons, respectively. The lowest CV recorded in *kharif* (33.0%) indicates more reliable receipt of rainfall. The rainy days almost followed the same trend as that of rainfall (Table 1).

Monthly rainfall

The data on monthly rainfall (Table 1) revealed that August was the highest (321.6 mm) rainfall receiving month followed by July (318.9 mm). November was observed to be the least rainfall contributing month (0.7%). The period from June to August was the most dependable (CV range 36-55%) and the months of October to December were the most uncertain (CV range 162-185%).

Weekly rainfall

The weekly average rainfall ranges from 0.4 mm (44th SMW) to 90.3 mm (31st SMW). Mean weekly precipitation amount and its assurance was low during the initial 22 SMWs, reaches the peak (>50 mm/week)

during 27th SMW to 36th SMW and again declined thereafter. The weekly average rainfall during the monsoon season (June to September) is higher than other seasons varying from 15.7 mm to 90.3 mm (Fig. 1). The CV values during monsoon season were relatively less (69–148%) owing to stable quantum of rainfall compared to the values of 131–456 per cent and 145–297 per cent in post and pre-monsoon weeks, respectively (Table 2). The higher values of CV in all weeks indicated that the rainfall was much erratic in distribution and it was higher in post monsoon weeks.

Probability of dry and wet spells

The probability of occurrence of dry spells (Table 2) is as high as 80-100 per cent in the first 22 SMWs except for 8th SMW where it was only 76 per cent of getting a dry week. Thereafter, the values remain low between 26th to 33rd SMW (representing wet spell sequence) and then again continue to increase till the end of the year. On the other hand, the probability of getting rainfall (wet spell) remains very high (71-86%) during 26th to 35th SMW except for 34th SMW. The probability of getting a wet spell, two and three consecutive wet spells are low during the first 25 SMWs. The values remained high during 26th to 35th SMW and again very low from 36th SMW till the end

Table 2: Mean weekly rainfall, its variability and probability of different dry and wet spells at Ballawal Saunkhri (1984-2004)

MSW	Rainfall (mm)	p(d)	p(dd)	p(ddd)	p(w)	p(ww)	p(www)
1	8.8	86	71	52	14	0	0
2	9.6	81	67	67	19	5	0
3	8.6	86	67	67	14	5	0
4	5.4	95	81	71	5	5	0
5	6.6	86	86	52	14	5	0
6	9.1	81	71	52	19	0	0
7	10.9	81	57	57	19	0	0
8	14.7	76	62	62	24	5	0
9	7.0	86	71	71	14	0	0
10	7.0	95	81	81	5	0	0
11	11.3	100	90	76	0	0	0
12	7.0	90	90	71	10	0	0
13	6.1	90	81	71	10	0	0
14	4.6	90	81	76	10	0	0
15	6.8	86	81	76	14	0	0
16	1.1	100	86	76	0	0	0
17	4.7	95	86	81	5	0	0
18	6.4	81	81	81	19	10	0
19	8.1	90	86	76	10	0	0
20	6.2	90	76	76	10	0	0
21	9.2	95	81	57	5	0	0
22	7.1	86	81	38	14	10	0
23	15.7	67	62	24	33	5	0
24	18.6	76	38	38	24	19	10
25	20.7	62	48	24	38	29	10
26	39.9	29	19	19	71	43	38
27	50.9	29	0	5	71	52	43
28	78.5	29	5	0	71	67	52
29	65.7	19	10	0	81	62	57
30	74.9	14	10	0	86	67	57
31	90.3	19	0	0	81	71	62
32	83.1	19	5	0	81	67	48
33	78.9	14	0	0	86	62	48
34	50.2	33	5	0	67	52	38
35	62.2	19	5	0	81	52	14
36	68.9	38	14	0	62	19	0
37	18.4	67	24	10	33	5	0
38	43.6	62	33	10	38	5	0
39	17.4	76	57	24	24	19	0
40	6.7	86	67	67	14	5	0
41	11.9	86	81	67	14	0	0
42	15.2	90	67	67	10	0	0
43	0.8	100	81	81	0	0	0
44	0.4	100	100	100	0	0	0
45	1.9	95	95	86	5	0	0
46	0.6	100	95	86	0	0	0
47	13.8	86	81	81	14	0	0
48	0.8	100	86	86	0	0	0
49	8.1	95	95	81	5	0	0
50	2.0	100	95	71	0	0	0
51	10.7	86	81	76	14	0	0
52	8.4	86	71	67	14	0	0

of the year. The probability of getting a month being dry remains in the range of 85 – 100 per cent for the entire pre and post monsoon season months (Fig. 2). However, the months of July and August were found to have 95–100 per cent assured rainfall followed by September (76 %) and June (48%).

Crop planning

Based on the above analysis, the following recommendations for the region could be made to increase the crop production per unit area under rainfed conditions. About 80 per cent of the total average annual rainfall coincides with *kharif* season and is received during a short time span of two and half months between July to September due to south-west monsoon. Rainfall received during April–May can be utilized for summer ploughing to make the land ready for final field preparation. The earliest start of rainy season had occurred in 24th SMW (June 11-17) and delay in start was as late as 29th SMW (July 16-22). The normal start of rainy season was observed as 26th SMW (June 25-July 1) in the region with coefficient of variation of 5.2 per cent. With the normal onset of rainfall, sowing of main season *kharif* crop like maize (PMH 2, JH 3459) should be started from the second fortnight of June. In case of delayed start, short duration drought tolerant maize varieties (megha, parkash) and low water requiring crops like oilseeds (sesame-TC 289, Punjab Til No. 1) and pulses (greengram-SML 668, ML 613, ML 818 / blackgram- mash 338) should be grown accordingly. In the event of mid season drought, mulching will help in reducing soil evaporation and conserving moisture in top layers of the soil. In the event of terminal drought, receding soil moisture conditions, crop requires supplementary irrigation. Intercropping of maize (50/60cm) with pulses (greengram/blackgram) in 1:1 row proportion can be viable option for increasing per unit area crop productivity under rainfed conditions.

A major portion of monsoon rainfall is generally lost through runoff (30-40%), which can be stored

through the construction of suitable water harvesting structures as on-farm reservoirs. The rainfall received during October-November is only 4-5 per cent of the total average annual rainfall which is very low for the sowing of *rabi* season crops. Therefore, soil and moisture conservation measures need due attention to conserve rainwater particularly during the months of July to September. Sowing of main *rabi* season crops like wheat (PBW 175 and PBW 527) and mustard (PBR 97, RLM 619) should be started from the last week of October. In case, the timely sowing of *rabi* crops is not done due to inadequate soil moisture, then wheat varieties like PBW 373 and PBW 509 and mustard like RLM 619 could be sown with the receipt of late winter showers for taking remunerative crop yields.

REFERENCES

- Dingre, S. and Habib, R. (2006). Rainfall analysis and crop planning for Kashmir valley. *J. Agrometeorol*, 8(2): 281-286.
- Jadhav, J.D., Mokashi, D.D., Shewale, M.B. and Patil, J.D. (1999). Rainfall probability analysis for crop planning in scarcity zone of Maharashtra. *J. Agrometeorol*, 1(1): 59-64.
- Robertson, G.M. (1976). Dry and wet spell UNDP/FAO, Tan Razak Agric. Res. Center, Sungh: Tekam, Malaysia, Project Field Report, *Agrometeorol. A-6*: 15.
- Sarma, N.N., Paul, S.R. and Sharma, D. (1996). Rainfall pattern and rainfall based cropping system for hills zone of Assam. *Ann. Agric. Res.* 17(3): 223-229.
- Subramaniam, A.R. and Raju, P.A.N. (1988). Rainfall variability and crop production in North Coastal Andhra. *Fert. News*, 33(4): 39-49.
- Tiwari, A.K., Sharma, A.K. and Srivastava, M.M. (1992). Probability analysis of rainfall data of Datia district, Bundelkhand for crop planning, *Indian J. Soil Cons.*, 20(3): 82-88.