Biomass production of soybean in relation to evapotranspiration and temperature

AJAY K. SRIVASTAVA 1, A.S.R.A.S. SASTRI AND B.D. GUPTA

Department of Physics and Agrometeorology Indira Gandhi Agricultural University, Raipur-492012.

ABSTRACT

Soybean is a fast upcoming crop in monsoon season after rice in Chhattisgarh area of central India under rainfed conditions. To find out the relationship between evapotraspiration (ET), temperature and biomass production, experiments were conducted in monsoon seasons of years 1994 and 1995. Biomass production of soybean in relation to cumulative ET and accumulated heat units has been evaluated and expressed as linear regression.

Key words: Evapotraspiration, Growing dergee days, Biomass production, Water balance

In the Chhattisgarh region under rainfed condition farmers usually take rice (about 3.8 million hectares) in bunded fields and small millets and pigeonpea in unbunded fields. Due to higher requirements of water for rice and decreasing trends of rainfall, a recommendation has been made for soybean in unbunded black soils under rainfed condition in monsoon season. In the recent decade, since that recommendation, the area under soybean has increased from 3,000ha to more than 85,000ha in monsoon season in the Chhattisgarh region.

The amount of ET loss for a given crop depends on climatic parameters, type of soils and growth stage of crop and crop types. Allison et.al. (1958) and Hanks et.al. (1968) studied the relationship between ET and biomass production of different crops grown in lysimeters under rainfed condition. They reported that a linear relationship becomes

valid after sufficient ET has occurred from crop establishment in initial stages.

Doorenbos and Pruitt (1977) expressed that the productivity of a crop is also dependent on degree days and also the knowledge of water loss through plant and its utilization in biomass production is an important factor. Cumulative ET (Chakravarty and Sastry, 1983), and degree days (Uchijma, 1975) have been reported to be highly correlated with crop biomass production.

MATERIALS AND METHODS

The experiments were conducted in the years 1994 and 1995 at the research farm of Indira Gandhi Agricultural University, Raipur to examine the evapotranspiration (ET) and degree day relationship with biomass production of soybean (Cv. JS 72-44) during monsoon season under rainfed

Present Address: Division of Agricultural Physics, IARI, New Delhi -110012

condition. Two sets of gravimetric Evapotranspiration and rainfall lysimeters were used for ET measurement. Soybean was sown by seed drill on 3rd July in 1994 and 22nd June in the year 1995 with a row spacing of 30cm. N.P and K were given as locally recommended. The crop was harvested on 7th November in 1994 and 31st October in 1995, Growing degree days (GDD) were computed using a base temperature of 10 °C.

The meteorological data were collected from the experimental field. Above and below ground biomass samples were collected randomly from 5 points at weekly intervals. Leaf, stem, pod and root were dried separately at 70 °C for 72 hours to obtain a constant dry weight.

RESULTS AND DISCUSSION

Water balance

Weekly water balance has been computed from sowing to maturity following the Thornthwaite and Mather book-keeping procedure (1955) and shown in Table 1. The potential evapotraspiration (PET) values required for the water balance computations are estimated by using Penman's equation (1948). From the Table 1 it can been seen that the in grain filling stage (38-41 SMW) the water deficit is more in the year 1995 as compared to the same stage of 1994.

There was water stress during mid seedling and end of reproductive stages of soybean crop in 1995 but there was very little water stress during the end of reproductive stage in 1994 (Srivastava et.al., 1996). As a result of this, the productivity of soybean decreased from 2.3 t had to 1.7 t had (a decrease of 0.55 t hard). Total biomass during 1994 was 5.51 t hard as compared to 6.13 t harduring 1995.

The evapotranspiration (ET), Pan evaporation (EP), rainfall (R) during different growth stages for the two kharif seasons are shown in Table 2. The total AET(lysimeter) losses during the growing season of soybean varied from 562.2 mm in 1994 to 525.7mm in kharif 1995. The EP values varied from 433.9 mm in 1994 to 523.0 mm in 1995 indicating that the AET and EP values varied independently between the seasons in the different phases.

In the two years of experiment, crop experienced two different rainfall distributions at Raipur. During 1994 rainfall was 1159.4mm and 1995 it was 927.4mm. Rainfall and soil moisture (SMW 30-33) have affected the AET(lysimeter) rate in 1994 and the weekly AET(lysimeter) value reached 45.8mm in vegetative stage which is more compared to the same stage AET (lysimeter) in 1995. This indicates that in the year 1994 the soybean crop AET(lysimeter) was much higher than the corresponding phenophase of 1995 (Table 2).

Crop stagewise ET, biomass and GDD

The stagewise crop AET (lysimeter), biomass and GDD are shown in Table 2 and 3. The AET (lysimeter) losses are highest during the vegetative and reproductive stages accounting more than 80% of the total AET where as the GDD accounting more than 65% of its total during these two stages. In the year 1994 the average dry biomass production was 5510 kg/ha as compare to 6192 kg/ha in 1995(+11.3 %).

Evapotranspiration and biomass production

The patterns of cumulative AET

Table 1: Weekly Climatic water balance (Thornthwaite method) at Raipur during kharif 1994 and 1995

		1994					1995					
SMW*	Rain (mm)	PET (mm)	ET (mm)	WS (mm)	WD (mm)	MAI (mm)	Rain (mm)	PET (mm)	ET (mm)	WS (mm)	WD (mm)	MAI (mm
25	-	-	-	-			22.2	12.6	12.6	0.0	0.0	1.0
26	- 3	-	-	-	0€	*	8.6	38.7	15.4	0.0	23.3	0.4
27	71.2	29.1	29,1	36.1	0.0	1.0	41.2	35.7	35.7	0.0	0.0	1.0
28	258.4	18.4	18.4	240.0	0.0	1.0	74.0	25.3	25.3	0.0	0.0	1.0
29	184.2	19.6	19.6	164.6	0.0	1.0	150.6	19.1	19.1	0.0	0.0	1.0
30	15.8	26.2	26.0	0.0	0.2	0.99	141.2	25.5	25.5	27.5	0.0	1.0
31	50.2	20.2	20.2	19.8	0.0	1.0,		24.2	24.2	37.6	0.0	1.0
32	55.4	20.2	20.2	35.2	0.0	1.0	78.8	24.6	24.6	54.2	0.0	1.0
33	75.2	25.2	25.2	50.0	0.0	1.0		26.9	26.9	124.1	0.0	1.0
34	112.6	24.1	24.1	88.5	0.0	1.0	16.4	27.1	26.9	0.0	0.0	0.99
35	111.0	21.1	21.1	89.9	0.0	1.0	111.2	25.5	25.5	75.2	0.0	1.0
36	111.8	25.8	25.8	86.0	0.0	1.0	4.8	28.0	27.3	0.0	0.7	0.97
37	21.8	27.3	27.3	0.0	0.1	1.0	3.0	26.8	24.5	0.0	2.3	0.92
38	10.8	24.5	23.8	0.0	0.7	0.97	21.2	29.0	27.9	0.0	12.1	0.96
39	39.2	26.1	26.1	0.0	0.0	1.0	13.0	28.2	25.7	0.0	2.5	0.91
40	4.4	25.8	24.5	0.0	1.3	0.95	0.0	30.4	23.8	0.0	6.5	0.78
41	31.0	24.1	24.1	0.0	0.0	1.0	2.6	26.0	19.6	0.0	6.4	0.75
42	2.0	25.3	22.6	0.0	2.7	0.89	25.8	24.0	24.0	0.0	0.0	1.0
43	1.8	23.1	19.0	0.0	4.1	0.82	0.0	22.6	15.5	0.0	7.1	0.68
44	2.6	18.1	14.2	0.0	3.9	0.79	0.0	9.6	**	0.0	3.4	0.64
	1159.4	424.2	411.3	810.1	13.0	0.00000	927.4	509.8	450.0	318.6	64.3	0.04

Where WS = Water surplus
WD= Water deficit
MAI = Moisture availability index
WS = Water surplus
Seedling
Vegetative
Vegetative
Reproductive
30-35
Reproductive
36-41

(lysimeter) and biomass production at different days after sowing (DAS) are shown in Table 4. From the table it can been seen that in both the years there is variations in AET(lysimeter) and biomass production. In the year 1994 cumulative AET (lysimeter) is high than the year 1995 but the biomass production is less in 1994 as compared to 1995. To find out the relationship between the AET (lysimeter) and biomass production correlation coefficients and regression analysis are worked out and presented below:

The correlation and regression equation between cumulative AET (lysimeter) and biomass production was worked out and is given below:

SMW

42-44

Correlation coefficient	Regression equation
0.94**	Y=3.4X-553

** Significant at 1% level

Y=Biomass production in grams meter2

X=Cumulative AET(lysimeter) in mm

Table 2: Stagewise AET (lysimeter) and rainfall of soybean crop at Raipur in Kharif season.

CROP STAGE		1994			1995	
	AET(mm)	EP(mm)	Rain(mm)	AET(mm)	EP(mm)	Rain(mm)
Seedling (3)*	52.2	75.4	510.0	35.0	109.3	72.0
Vegetative(6)	242.5	140.8	421.2	181.3	140.6	657.4
Reproductive(6)	238.1	148.2	218.2	247.7	166.2	169.6
Maturity(3)	29.4	69.5	6.6	61.7	107.0	28.4
Total (18)	562.2	433.9	1156.0	525.7	523.1	927.4

^{*} Figures in brackets are weeks

Table 3: Stagewise GDD and biomass in two years of experiments.

Crop stage	GDD(°C) Total	1994 Biomass(Kg/ha)	GDD(°C)	1995 Total Biomass(Kg/ha)
Seedling(3)*	458	150	341	180
Vegetative(6)	660	2054	744	2490
Reproductive(6)	676	3276	752	3502
Maturity(3)	222	30	352	20
Total	2016	5510	2189	6192

Figures in brackets are weeks

Table 4: Cumulative AET (lysimeter) and total dry matter production of soybean crop.in the year 1994 and 1995.

Days after	Cumulative	eET(mm)	Biomass production(gm ⁻²)		
sowing(DAS)	1994	1995	1994	1995	
32	132	85	17	33	
57	219	152	45	83	
520	284	239	182	251	
62 72	327	303	438	445	
82	396	364	800	686	
92	460	421	1121	960	
102	519	466	1178	1252	
112	543	486	1332	1470	
126	566	528	1339	1482	

The correlation was highly significant in both the years. The total AET (lysimeter) requirement for soybean crop varied between 500-550mm in Raipur climatic condition and it varied accordingly with rainfall amount and its pattern of distribution in crop growing season.

GDD and biomass production

At different accumulated GDD levels, biomass production in two years are shown in Table 5. In the year 1995 during earlier phenophases there was more accumulation of biomass as compared to the year 1994 for same value of GDD. But at later stages in the year 1995 the rate of accumulation of biomass is more as compared to the year 1994. The GDD calculation has been carried out upto 1700 GDD i.e. starting of physiological maturity.

Table 5: Biomass production of soybean crop at different unit of GDD

GDD(°C)	Biomass Production (g		
	1994	1995	
400	17	26	
600	40	63	
900	285	222	
1000	438	445	
1300	735	608	
1500	1056	915	
1700	1150	1205	

The correlation and regression equation between growing degree days and biomass production of soybean pooled for two years is given below:

Y = 1.2GDD - 598 (r=0.96**)

** Singnificant at 1% level

Y = Biomass production in grams meter²

GDD= Growing degree days in °C

ACKNOWLEDGEMENT

The authors are thankful to the Indian Council of Agricultural Research for the financial support and providing a Project "Water balance studies of Soybean - Chickpea sequential cropping system under rainfed condition in Agroclimatic Zone VII".

REFERENCES

Allison, L.H., Roller, E.M. and Raney, W.A. 1958. Relationship between evapotranspiration and yield of crop grown in lysimeters receiving natural rainfall. Agronomy Journal, 50:506-511.

Chakravarty, N.V.K. and Sastry, P.S.N. 1983. Biomass production in wheat in relation to evaporative demand and ambient temperature. Mausam, 34(3):323-326.

Dorrenbos, J. and Pruitt, W.V. 1977. Crop water requirement. FAO irrigation and drainage paper no-24 FAO, Rome pp 21-23.

Hanks, R.J., Gardener, H.R. and Florain, R.L. 1968. Evapo-transpiration-Climatic relationship for several crops in central great plains. Agronomy Journal, 60:538-592.

Penman H.L. 1948. Natural evaporation from open water, bare soil and grass. Proceeding of Royal Society London (Sec. A), 193:120-145.

Srivastava, A.K., Naidu, D., Sastri, A.S.R.A.S., Urkurkar, J.S. and Gupta, B.D. 1996. Effects of water stress on soybean productivity in central India, Drought Network News, 8(1):7-8.

Thronthwaite, C.W. and Mather, J.R. 1955.
The water balance In Climatology,
John Hopkins, USA, University Lab of
Climatology, New Jersey

Uchijma,Z. 1975. Dry matter production of crops in relation to climatic conditionscollected papers of Agriculture Meteorology No16. National Institute of Agricultural Science Japan, 2-5.