Short communication

Estimating crop water requirements for irrigation scheduling in different crops in humid subtropical agro-climate of Western Himalayas

ARUNAVA PODDAR1*, VIJAY SHANKAR2 and NAVSAL KUMAR1

¹Dept of Civil Engineering, Shoolini University, Solan (HP), India ²Dept of Civil Engineering, NIT Hamirpur (HP), India *Corresponding Author: arunava.nithrs@gmail.com

Deficiency of water in the rhizospheric area of soil results in reduced crop growth adversely affecting crop yield. Thus, the objective of irrigation is to maintain the adequate moisture content in the root zone, such that the crop yield is not adversely affected (Kumar et al. 2012; Kumar et al. 2020). However, in the current scenario of climate change, rapid industrialization and population increase, there is tremendous pressure on water resources both quantitatively and qualitatively (Rao and Poonia 2011). Hence, precise allocation of water resources considering the crop water requirement and proper knowledge of soil composition is essential to attain optimum yield, and maximum water use efficiency (Mehta and Pandey 2016). The crop water requirement is generally crop evapotranspiration (ET₂) considering atmospheric water loss through plant transpiration and soil evaporation simultaneously (Kumar 2017; Poddar et al. 2021). There are several methods for direct estimation of ET which includes energy balance, microclimatological methods, field water balance and Lysimeters. However, the indirect methods includes measurement of reference evapotranspiration and crop coefficients (Chaudhari et al. 1999). Estimating water balance in the Lysimeter is the most reliable and useful approach for determining actual ET under field conditions (Kashyap and Panda, 2001). Hence, the present study was undertaken for rabi (Wheat, Indian mustard, Potato) and kharif (Maize, Sorghum, Guar) crops grown in a humid subtropical agro-climate of western Himalayas with a specific objective to estimate the ET_c using the lysimeter water balance approach.

Description of the field experiments

An experiment was conducted in the campus of National Institute of Technology Hamirpur, Himachal Pradesh, India situated at 3162.8'68". N latitude and 7613.3'52". E longitude, and 895 m amsl altitude. The

field trials were conducted during the 2017 – 2019 crop growing seasons. The crops considered in the experimental study were wheat (*Triticum aestivum*), Indian mustard (*Brassica Juncea*), potato (*Solanum tuberosum L.*), maize (*Zea mays*), sorghum (*Sorghum bicolor L. Moench*) and guar (*Cyamopsistetragonoloba L.*). The details of the crop type, crop duration, growth stages, and irrigation events for the present study are illustrated in Table 1. Local agronomic practices (i.e., land preparation, manuring, fertilization) were followed while conducting experiments on different crops.

Water balance method

Two drainage type lysimeters $(1.5\times1.5\times2 \text{ m})$ were installed separately and rim of the lysimeter was kept 0.1 m above ground level to prevent surface runoff. A land gravel filter (0.3m) was provided at the bottom of the lysimeter to facilitate drainage to a calibrated collector. A soil moisture capacitance probe was used to measure soil moisture content at 0.1 m interval till 1.6 m in the soil profile on daily basis. The applied irrigation was measured using a discharge meter installed at the outlet of the water tank. The daily rainfall was recorded using a tipping bucket rainfall.

The water balance method assesses the incoming and outgoing water flux into the crop root zone. Since drainage type lysimeters were used, ET_c was obtained for long periods. The ET_c was determined using the water balance equation (Bandyopadhyay and Mallick, 2003) as:

$$P + I = D + ETc + RO \pm \Delta S$$
 (1)

where, P = precipitation (mm), I = irrigation (mm), D = Drainage from lysimeter (mm), RO = runoff (mm), and ΔS = change in soil moisture storage (mm). ΔS for a specific depth (d_z) for a specific time period is computed as:

Table 1: Details of the crop duration, growth stages, and irrigation days pertaining to *Rabi* and *Kharif* crops grown in the field

Crop	Variety	Date of sowing	Date of harvesting	Duration	Growth stages (Days)				Irrigation	Spacing
					I	II	III	IV	provided (DAS)	(cm)
Rabi Crops										
Wheat (Triticum aestivum)	Super (6776/PB)	3 rd Jan,2017	15 th May, 2017	133	25	36	45	28	$26^{\text{th}},44^{\text{th}},56^{\text{th}},\!80^{\text{th}},\\96^{\text{th}},116^{\text{th}}$	20x5
Indian mustard (Brassica juncea)	VL-804	22 nd Jan, 2018	14 th May, 2018	113	19	32	38	25	11 th , 25 th , 37 th , 59 th , 91 st	40x15
Potato (Solanum tuberosum L.)	Kufri Himsona	7 th Jan, 2019	6 th May, 2019	120	22	32	38	29	21 st ,40 th , 52 nd , 64 th ,87 th , 104 th	45x15
Kharif Crops	-									
Maize (Zea mays)	Maize 7074 (HYBRID)	20 th May, 2017	10 th Sept, 2017	114	20	34	36	24	22 nd ,36 th , 48 th , 64 th	50x20
Sorghum (Sorghum bicolor L. Moench)	TX 610	16 th May, 2018	22 nd Sept, 2018	130	21	35	39	35	$22^{nd},48^{th},75^{th},\\93^{rd}$	30x15
Guar (Cyamopsistetr- agonoloba L.)	Agaita Guara - 112	28 th May 2019	1st Sept, 2019	97	20	26	28	23	$26^{th}, 53^{rd}, 81^{st}$	25x15

$$(\Delta Sz) = (\theta z, \text{final} - \theta z, \text{initial}) \times dz$$
 (2)

where $\theta_{z, \; final}$ and $\theta_{z, \; initial}$ is the final and initial moisture content, respectively in the soil profile in a discrete-time interval.

ET variation in rabi crops

The seasonal ET_c of rabi crops, i.e., Indian mustard, wheat, and potato, was 165.8 mm, 242.7 mm, and 308.7 mm, respectively (Table 2). The precipitation received during the crop period of wheat, Indian mustard, and potato was 112.4 mm, 114.5 mm, and 108.5 mm, respectively. The amount of irrigation required for wheat, Indian mustard, and potato, was 195.0, 140.0, and 240.0 mm, respectively. This indicated that the irrigation requirements of rabi crops are higher than the precipitation received during the crop period. In the case of wheat and potato, the irrigation supplied was almost twice the precipitation, indicating that nearly twothirds of the crop water requirements were fulfilled by irrigation. The contribution to the groundwater during the rabi crop season was quite low as compared to the amount of water received. The variation of stage-wise ET for rabi crops considered has been shown in Table. 2. It is evident from the table that the ET_c during the initial and midseason stage for rabi crops was quite similar. However,

during the crop development stage, the $\mathrm{ET_c}$ for potato was significantly higher than wheat and Indian mustard. During the late season, the $\mathrm{ET_C}$ for wheat was higher as compared to the other two crops.

ET variation in kharif crops

The seasonal ET of kharif crops, i.e., maize, sorghum, and guar, was 502.9 mm, 518.9 mm, and 494.7 mm, respectively (Table 2). The amount of precipitation obtained during the crop period of maize, sorghum, and gaur, was 1052.5, 1059.3, and 1124.3 mm, respectively, which is comparatively much higher than the rabi crops. The total amount of irrigation required for maize, sorghum, and gaur, were 10.0, 31.4, and 20.0 mm, respectively, which is substantially less than the irrigation required for rabi crops. This also indicates that the irrigation requirements of kharif crops were very low as compared to the received precipitation. More than 95 per cent of the crop water requirements were fulfilled by the precipitation alone. The groundwater contribution during the *kharif* crop season was substantially high. The variation of stage-wise ET_c for kharif crops considered has been shown in Table 2. The ET for kharif crops during the initial and late-season stages was nearly the same. The ET_c for sorghum was significantly low during the crop

Table 2: Crop growth stage-wise water balance components for *Rabi* and *Kharif* crops

Water balance component	Initial	Development	Mid- season	Late- season	Total (mm)	Initial	Development	Mid- season	Late- season	Total (mm)		
Wheat						Indian mustard						
P	0.0	6.8	104.6	1.0	112.4	6.0	10.5	11.5	77.0	105.0		
Ir	20.0	70.0	35.0	70.0	195.0	20.0	60.0	70.0	20.0	170.0		
Dr	8.9	17.5	39.5	4.0	69.9	6.2	7.9	5.4	51.1	70.6		
ΔS	-7.6	21.2	-10.5	-8.3	-5.2	-1.6	4.7	-9.7	24.4	17.8		
$\mathrm{ET_{c}}$	18.7	38.1	110.6	75.3	242.7	21.4	57.9	85.8	21.5	186.6		
Potato						Maize						
P	0.0	6.8	4.7	97.0	108.5	152.4	385.7	241.4	273	1052.5		
Ir	20.0	140.0	80.0	0.0	240.0	0.0	10.0	0.0	0.0	10.0		
Dr	4.5	6.0	5.4	42.0	57.9	101.2	176.4	88.4	187.6	553.6		
ΔS	-10.6	-11.4	-21.7	15.6	-28.1	30.0	-16.4	-18.3	10.7	6.0		
ET_{c}	26.1	152.2	101.0	39.4	318.7	21.2	235.7	171.3	74.7	502.9		
Sorghum						Guar						
P	61.5	203.5	552.8	241.5	1059.3	132.6	485.7	280.4	225.6	1124.3		
Ir	31.1	0.0	0.0	0.0	31.1	20.0	0.0	0.0	0.0	20.0		
Dr	39.2	96.2	284.2	129.5	549.1	95.2	252.3	165.2	161.5	674.2		
ΔS	-2.0	12.3	10.2	1.8	22.3	25.0	8.4	-29.3	-28.7	-24.6		
ET _c	55.4	95.0	258.3	110.2	519.0	32.4	225.0	144.5	92.8	494.7		

Note: P - Precipitation, I_r -Irrigation, D_r - Percolation to groundwater, ΔS - Change in soil moisture storage, and ET_c -Crop evapotranspiration

development and significantly high during the mid-season stage when compared to maize and guar crops. The $\mathrm{ET_c}$ variation of maize and guar crops was similar during all stages of crop growth.

The seasonal ET_c in the case of *rabi* crops was highly variable, but for *kharif* crops, it was nearly similar. The stage-wise ET_c variation was different for each *rabi* crop considered. For *kharif* crops, maize and guar followed similar stage-wise ET_c, whereas sorghum followed a different pattern. The estimated ET_c of different crops may further help in planning of optimal irrigation schedules in the humid sub-tropical agro-climate of the western Himalayas.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the financial support received from the Ministry of Earth Sciences, Govt. of India.

Conflict of Interest Statement : The author(s) declare(s) that there is no conflict of interest.

Disclaimer: The contents, opinions, and views expressed in the research article published in the Journal of Agrometeorology are the views of the authors and do not necessarily reflect the views of the organizations they belong to.

Publisher's Note: The periodical remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

REFERENCES

Bandyopadhyay, P. K. and Mallick, S. (2003). Actual evapotranspiration and crop coefficients of wheat (*triticuma estivum*) under varying moisture levels of humid tropical canal command area. *Agric. Water Manage.*, 59: 33-47.

- Chaudhari, G. B., Patel, K. I., Shekh, A. M. and Savani, M. B. (1999). Crop coefficients of major crops in middle Gujarat region. *J. Agrometeorol.*, 1(2): 167-172.
- Kashyap, P. S. and Panda, R. K. (2001). Evaluation of evapotranspiration estimation methods and development of crop-coefficients for potato crop in a sub-humid region. *Agric. Water Manage.*, 50(1): 9-25.
- Kumar, N., Poddar, A., Shankar, V., Ojha, C. S. P. and Adeloye, A. J. (2020). Crop water stress index for scheduling irrigation of Indian mustard (*Brassica juncea*) based on water use efficiency considerations. *J. Agron. Crop Sci.*, 206(1): 148-159.
- Kumar, R., Jat, M. K. and Shankar, V. (2012). Methods to estimate reference crop evapotranspiration- A

- review. Water Sci. Technol., 66 (3): 525-535.
- Kumar, S. (2017). Reference evapotranspiration (ET₀) and irrigation water requirement of different crops in Bihar. *J. Agrometeorol.*, 19(3): 238-241.
- Mehta, R., and Pandey, V. (2016). Crop water requirement (ET_c) of different crops of middle Gujarat. *J. Agrometeorol.*, 18(1): 83.
- Poddar, A., Gupta, P., Kumar, N., Shankar, V. and Ojha, C.S. (2021). Evaluation of reference evapotranspiration methods and sensitivity analysis of climatic parameters for sub-humid sub-tropical locations in western Himalayas (India). *ISH J Hyd. Engg.*, 1-11. https://doi.org/10.1080/09715010.201 8.155173
- Rao, A. S., and Poonia, S. (2011). Climate change impact on crop water requirements in arid Rajasthan. *J. Agrometeorol.*, 13(1): 17-24.

Received: May 2020: Accepted: May 2021